

**Gentamicin, Amphotericin B (GA)  
Antimicrobial Supplement  
Specification and Instruction Sheet**



**Quality Control Testing**

- Sterility
- Cell Culture Evaluation

| Product   | Part No.                | Volume | Concentration of Supplement                    | Storage |
|---|-------------------------|--------|--|---------|
| <b>Gentamicin, Amphotericin B (GA)<br/>Antimicrobial Supplement</b> | <a href="#">LS-1104</a> | 0.5 mL | Gentamicin 30 mg/mL<br>Amphotericin B 15 µg/mL | -20°C   |

**Gentamicin and Amphotericin B**

The use of GA is recommended to inhibit potential fungal or bacterial contamination of eukaryotic cell cultures. A complementary vial of GA is provided for your convenience.

Addition of 0.5 mL of GA to 500 mL of media will provide an effective concentration to inhibit bacterial and fungal cell division.

**Product Storage**

GA is best stored at -20°C until ready to use.

GA should only be thawed once and stored at 4°C for a maximum of two weeks after thaw. Multiple freeze/thaw cycles are not recommended.

Do not use product beyond expiration date.

**Safety and Use Statement**

GA Antimicrobial Supplement is for Research Use Only. This product is not approved for human or veterinary use or for use in *in vitro* diagnostics or clinical procedures.

To place an order, please visit [lifelinecelltech.com](http://lifelinecelltech.com) or call technical and customer service at 877.845.7787.

## Basic Aseptic Technique

GA Antimicrobial supplement, cells, medium, and reagents should only be used in an aseptic environment, a Class II biological safety cabinet with front access and filtered laminar airflow, or an equivalent device. Always wear gloves and eye protection when working with these materials.

Wipe or spray all bottles and vials with 70% ethanol or isopropanol, especially around the area of the cap, before placing them in the biological safety cabinet. Allow these surfaces to dry completely before opening the bottle or vials.

Transfer cells, medium or reagents with disposable sterile pipettes. Do not mouth pipette! Take up the volume needed into the pipette, being careful not to touch the sterile tip to the rim of the container or any other surface. Close the container and open the container into which the transfer is being made, again being careful not to touch any surfaces with the sterile tip. Transfer the material and close the container.

Wash your hands before and after working with cell cultures. Do not block airflow in a laminar flow hood as this may compromise sterility. Ensure that biological safety cabinets are certified routinely and that the HEPA filters are replaced regularly.

## The Lifeline Guarantee

Lifeline's rigorous quality control ensures sterility and performance to standardized testing criteria. Upon request, Lifeline will provide lot-specific QC test results, material safety data sheets and certificates of analysis. See complete guarantee/warranty statement at [lifelinecelltech.com](http://lifelinecelltech.com) or contact your Lifeline technical representative for more information.

**For any question on medium supplementation guidelines;  
please contact technical service at 877.845.7787.  
We are here to help.**

## Notes:

***Call Lifeline Technical Service and Sales at 877.845.7787  
or visit [lifelinecelltech.com](http://lifelinecelltech.com) for more information***

**Lifeline Cell Technology – 8415 Progress Drive, Suite T – Frederick, MD 21701**

©2016 Lifeline Cell Technology. All Rights Reserved